

EMC 451

Advanced ECG Interpretation

Unit 8: Preexcitation Syndromes

EMC 451: Preexcitation Syndromes

1

Unit Objectives

- Upon completion of this unit, you should be able to:
 - Define preexcitation.
 - Describe an accessory pathway.
 - Discuss why accessory pathways predispose patients to dysrhythmias.
 - Recognize the ECG characteristics of Lown-Ganong-Levine Syndrome (LGL) and Wolf-Parkinson-White Syndrome (WPW).
 - List the treatment of WPW and LGL.

EMC 451: Preexcitation Syndromes

2

Preexcitation

- Normally, conduction between the atria and ventricles is delayed about 0.1 seconds in the AV node.
- AV delay allows atria to contract and fill the ventricles.
- In preexcitation syndromes, an accessory pathway bypasses the AV node, permitting the ventricles to contract earlier than normal.

EMC 451: Preexcitation Syndromes

3

Accessory Pathways

- Several have been identified.
 - Kent bundle (WPW)
 - Intranodal bypass tract (LGL)
 - Mahaim fibers
 - Atriofascicular bypass tract
- Present in less than 1% of population.
- Occurs predominantly in males.

EMC 451: Preexcitation Syndromes

4

Wolf-Parkinson-White Syndrome


- Accessory pathway results from malformation of fibrous tissue that normally separates the atria and ventricles.
- Tissue provides a connection between atria and ventricles outside the conduction system (short-circuit).
- This tissue may provide a mechanism of AV reentry, thus the tendency to develop PSVT.

EMC 451: Preexcitation Syndromes

5

ECG Criteria of Wolf-Parkinson-White Syndrome


- PR interval < 0.12 sec.
- QRS complex > 0.10 seconds (20%)
- Delta wave (positive or negative)
- Secondary T-wave changes may be present
- Narrow complexes, no delta wave (80%)



EMC 451: Preexcitation Syndromes

6

Wolf-Parkinson-White Syndrome



Wolf-Parkinson-White Syndrome



EMC 451: Preexcitation Syndromes

8

Clinical Significance of Wolf-Parkinson-White Syndrome

- Atrial flutter (5%), atrial fibrillation (10-20%) and PSVT (40-80%) are the most frequently associated dysrhythmias
- Atrial fibrillation may be fatal because conduction over accessory pathway may result in ventricular rate of 200-300 bpm.
- Rapid ventricular rates may deteriorate into ventricular fibrillation.

EMC 451: Preexcitation Syndromes

9

Treatment of Wolf-Parkinson-White Syndrome

- If unstable, cardioversion.
- Vagal maneuvers
- Pharmacologically
 - Adenosine (narrow complex only)
 - Beta blockers (narrow complex only, but usually ineffective)
 - Verapamil (narrow complex only)
 - Other calcium channel blockers (narrow complex only)
 - Pronestyl (best treatment for wide complex)
 - Amiodorone
 - Flecainide
 - Propafenone
 - Sotalal
 - Do NOT use
 - Digoxin
 - Diltiazem
- Radio ablation

EMC 451: Preexcitation Syndromes

10

Long-Ganong-Levine Syndrome


- Accessory pathway bypasses the AV node, but conduction occurs via the usual ventricular conduction pathways.
- More common among women.
- Usually occurs between ages of 20 and 40.
- Shortened PR interval, but no delta wave.
- QRS is of normal width.

EMC 451: Preexcitation Syndromes

11

Long-Ganong-Levine Syndrome


Lown-Ganong-Levine (LGL) Syndrome



EMC 451: Preexcitation Syndromes

12

Long-Ganong-Levine Syndrome



EMC 451: Preexcitation Syndromes

13

Long-Ganong-Levine Syndrome

- Treatment is same as for WPW.
- Usually responds better than WPW and does not require ablation therapy.

EMC 451: Preexcitation Syndromes

14
